

Available online at www.sciencedirect.com

Discrete Mathematics 307 (2007) 748–749

DISCRETE
MATHEMATICSwww.elsevier.com/locate/disc

Note

Counterexample to a conjecture of Györi on C_{2l} -free bipartite graphs[☆]

C. Balbuena^a, P. García-Vázquez^b, X. Marcote^a, J.C. Valenzuela^c^a*Departament de Matemàtica Aplicada III, Universitat Politècnica de Catalunya, Campus Nord, Edifici C2, C/ Jordi Girona 1 i 3, E-08034 Barcelona, Spain*^b*Departamento de Matemática Aplicada I, Universidad de Sevilla, Avda Reina Mercedes 2, E-41012 Sevilla, Spain*^c*Departamento de Matemáticas, Universidad de Cádiz, Avda Ramón Puyol s/n, E-11202 Cádiz, Spain*

Received 24 November 2005; received in revised form 5 July 2006; accepted 10 July 2006

Available online 24 August 2006

Abstract

A counterexample on a conjecture of Györi related with C_{2l} -free bipartite graphs is described.
© 2006 Elsevier B.V. All rights reserved.

Keywords: Cycles in bipartite graphs; Forbidden subgraphs

In 1997 Györi [2] studied the structure of C_6 -free bipartite graphs and the relationship between this problem and some interesting results of Erdős et al. [1] on a number-theoretic problem. Namely, Györi proved a conjecture of Erdős et al. [1] regarding the maximum number of edges that a C_6 -free bipartite graph can have. Moreover, he proved another theorem that generalizes the previous one for cycles of longer length. In this paper Györi stated a conjecture [2, p. 373] that apparently contains a misprint¹ and it should have been expressed in this way:

Conjecture 1. If $G = (X, Y)$ is a bipartite graph with color classes X, Y where $|X| = m$, $|Y| = n$, $m^2 \leq n$, $3 \leq l \leq m$ and G has at least $(l-1)n + m - l + 2$ edges, then G must contain a cycle of length $2l$.


In a recent paper [3], the same author disproves Conjecture 1 for $l = 3$, but leaves the proof or refutation for $l \geq 4$ as an open problem. In this note we provide a counterexample that disproves Conjecture 1 when $m \geq 2l - 1$. Let us denote by $K_{(m,n)}$ the complete bipartite graph with m vertices in the first class and n vertices in the second one. Let us also denote by $d_G(v)$ the degree of the vertex v in the graph G .

Let l, m be integers such that $3 \leq l \leq 2(l-1) \leq m$. We consider the graphs $G_1 = K_{(m-l+1, l-1)}$ and $G_2 = K_{(l-1, n-l+2)}$. Take two vertices $u \in V(G_1)$ and $v \in V(G_2)$ with $d_{G_1}(u) = m - l + 1$ and $d_{G_2}(v) = l - 1$, and let G be the bipartite

[☆] Research supported by the Ministry of Education and Science, Spain, and the European Regional Development Fund (ERDF) under project MTM2005-08990-C02-02.

E-mail addresses: m.camino.balbuena@upc.edu (C. Balbuena), pgvazquez@us.es (P. García-Vázquez), francisco.javier.marcote@upc.edu (X. Marcote), jcarlos.valenzuela@uca.es (J.C. Valenzuela).

¹ The original conjectured value $(l-1)n + m - l + 1$ (see [2]) may be easily disproved by means of a graph roughly outlined by the author. We appreciate the referee's comments enlightening this misprint.

Fig. 1. The graph G with $l = 3$, $m = 5$ and $n = 25$.

graph on m and n vertices obtained by gluing the graphs G_1 and G_2 in such a way that the vertex u of G_1 is identified with the vertex v of G_2 (see Fig. 1).

Clearly, any cycle of G must be entirely contained in either G_1 or G_2 . But G_i , $i = 1, 2$, cannot contain a cycle of length $2l$ because one of its classes has cardinality $l - 1$. So G is free of C_{2l} and it has size $e(G) = (m - l + 1)(l - 1) + (l - 1)(n - l + 2) \geq (l - 1)n + m - l + 2$ because $m \geq 2l - 1$. Therefore, Conjecture 1 is disproved for $m \geq 2l - 1$.

In [2], Györi proved the following result:

Theorem. *If $G(X, Y)$ is a bipartite graph with color classes X, Y such that $|X| = m$, $|Y| = n$, $m^2 \leq n$ and G has at least $(l - 1)n + c(l)m^2$ edges for some constant $c(l)$ then G must contain a cycle of length $2l$.*

Thus, we propose the following reformulation of the conjecture:

Conjecture 2. *If $G = (X, Y)$ is a bipartite graph with color classes X, Y where $|X| = m$, $|Y| = n$, $m^2 \leq n$, $3 \leq l \leq m$ such that $m > (l - 1)^2$ and G has at least $(l - 1)n + 1/(l - 1)m^2$ edges then G must contain a cycle of length $2l$.*

Corollary of Theorem 1 in [3] confirms our Conjecture 2 for $l = 3$. For $l \geq 4$ it is still an open problem.

References

- [1] P. Erdős, A. Sarközy, V.T. Sós, On product representation of powers I, *European J. Combin.* 16 (1995) 567–588.
- [2] E. Györi, C_6 -free bipartite graphs and product representation of squares, *Discrete Math.* 165/166 (1997) 371–375.
- [3] E. Györi, Triangle-free hypergraphs, *Combinatorics, Probab. Comput.* 15 (1–2) (2006) 185–191.